



Common symptoms of obstructive HCM can include shortness of breath, rapid heartbeat, chest pain, feeling light-headed, fainting, and fatigue.

**“My obstructive HCM symptoms made me feel like I had to settle. But now my symptoms are improving, helping me go from expecting less to experiencing more.”**

THIS IS  
**MY CAMZYOS<sup>®</sup> MOMENT**

*Mike, a real CAMZYOS patient*  
Individual results may vary.

All people depicted are real patients and their care partners who were compensated for their time.

## Indication & Select Important Safety Information

### What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

### WARNING: RISK OF HEART FAILURE



**CAMZYOS may cause serious side effects, including heart failure.**

**Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.**

**You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS.**

Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • fatigue • leg swelling
- chest pain • racing heart (palpitations) • rapid weight gain

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

**The risk of heart failure is also increased when CAMZYOS is taken with certain other medications.** Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

**Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program.**

This means:

- ✓ You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.
- ✓ Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.
- ✓ You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.



If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.



# Introducing CAMZYOS

If you have symptomatic obstructive HCM, it may be helpful to learn about a different treatment option called CAMZYOS on the pages listed below.

What is symptomatic obstructive HCM? ..... 4–5

How CAMZYOS works ..... 6–7

Results from a clinical study with CAMZYOS ..... 9–13

Monitoring, safety measures, and precautions with CAMZYOS ..... 14–16

Meet patients on CAMZYOS ..... 17–18

Important Safety Information, including Boxed WARNING ..... 20–22

Side effects..... 22

MyCAMZYOS Patient Support program ..... 24

## Important Facts About CAMZYOS

### What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

### What should I discuss with my healthcare provider before starting CAMZYOS?



Talk to your healthcare provider about all of your medical conditions, including if:  
You are pregnant or plan to become pregnant—CAMZYOS may harm your unborn baby



Tell your healthcare provider right away if you become pregnant or think you are pregnant during treatment with CAMZYOS.  
You may also report your pregnancy by calling Bristol Myers Squibb at 1-800-721-5072 or [www.bms.com](http://www.bms.com).

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Learn more at [camzyos.com](http://camzyos.com)



CAMZYOS—THE FIRST AND ONLY FDA-APPROVED TREATMENT OF ITS KIND\*—TARGETS THE SOURCE OF SYMPTOMATIC OBSTRUCTIVE HCM.

\*CAMZYOS is a cardiac myosin inhibitor.





“I was starting to feel left out of my own life.  
I couldn’t join hikes with friends or family. Now, with CAMZYOS  
improving my symptoms, I’m hitting the trails with them again!”

*Stephanie, a real CAMZYOS patient*

Talk to your cardiologist today to see if a CAMZYOS  
moment may be in your future, too.

Individual results may vary.

Friends of Stephanie, shown here, are actor portrayals.

## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

## WARNING: RISK OF HEART FAILURE

**!** CAMZYOS may cause serious side effects, including **heart failure**.

Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS. Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.

- !** Get medical help right away if you experience new or worsening symptoms, including:
- shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

The risk of heart failure is also increased when CAMZYOS is taken with certain other medications. Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program. This means:

- ✓ You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.
- ✓ Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.
- ✓ You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.
- i** If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.

**!** CAMZYOS may cause serious side effects, including **heart failure** (a condition where the heart cannot pump with enough force).

- !** Get medical help right away if you experience new or worsening symptoms, including:
- shortness of breath • fatigue • leg swelling
  - chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.

- 💬** Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.

- 💬** Talk to your healthcare provider for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.




# What is obstructive HCM?

Obstructive HCM is the most common type of HCM—a condition that can affect the function of your heart.

HCM is a heart disease that can be inherited, which means it may run in your family due to genetics. It's estimated that HCM impacts between 1 in 200\* and 1 in 500† people.

HCM causes the heart muscle wall to thicken and stiffen, making it harder for the heart to fill with blood. When HCM causes the heart muscle to get so thick that it blocks or reduces blood flow from the heart to the rest of the body, it's called obstructive HCM.



The majority of people with HCM have been diagnosed with obstructive HCM.

\*The 2015 Semsarian publication identified that the prevalence of HCM gene carriers could be as high as 1 in 200.  
†The 1995 CARDIA study—a multicenter, US-population-based echocardiography study of 4111 subjects (aged 23–35)—identified the prevalence of HCM as 1 in 500 people in the general population.


Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Learn more at [camzyos.com](#)


# What are the symptoms of obstructive HCM?

Obstructive HCM makes it difficult for your heart muscle to relax, so your heart must work harder and use more energy.


Common symptoms can include:




Shortness of breath




Rapid heartbeat




Chest pain



Feeling light-headed



Fainting



Fatigue

If you have been diagnosed with obstructive HCM and are experiencing any of the symptoms above, talk to your doctor today.



# What is CAMZYOS?

If you have symptomatic obstructive HCM, talk to your doctor about CAMZYOS, a different treatment option that may be right for you.

CAMZYOS is a prescription medicine used to treat adults with symptomatic obstructive HCM and may improve both your symptoms and your ability to be active.



**CAMZYOS is a capsule that should be taken at the same time once a day.**

**It can be taken with or without food. Take CAMZYOS exactly as your healthcare provider directs; see the Medication Guide for more details.**

- Do not change your dose of CAMZYOS without talking to your doctor first
- Swallow the capsule whole without breaking, opening, or chewing it
- If you miss a dose of CAMZYOS, take it as soon as possible, and take your next dose at your regular time the next day. Do not take 2 doses on the same day to make up for a missed dose
- Your doctor may change, temporarily stop, or permanently stop your treatment with CAMZYOS if you have certain side effects
- If you take too much CAMZYOS, call your doctor or go to the nearest hospital emergency room right away

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Learn more at [camzyos.com](#)

## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

## WARNING: RISK OF HEART FAILURE



**CAMZYOS may cause serious side effects, including heart failure.**

Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS. Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

The risk of heart failure is also increased when CAMZYOS is taken with certain other medications. Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program. This means:



You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.



Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.



You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.



If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](#) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.



**CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).**



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • fatigue • leg swelling • chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.



**Tell your healthcare provider** if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.



**Talk to your healthcare provider** for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.


Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.



# How does CAMZYOS work differently?


CAMZYOS is the first and only FDA-approved cardiac myosin inhibitor. CAMZYOS targets the source of symptomatic obstructive HCM.

Normal heart




Two proteins in the heart muscle, called myosin and actin, contribute to your heart pumping by connecting (making the heart contract) and releasing (allowing the heart to relax).

Heart with obstructive HCM



In obstructive HCM, myosin and actin form too many connections with each other, causing the heart to contract too much, which can make it difficult for it to relax.

How CAMZYOS works in a heart with obstructive HCM



CAMZYOS binds to myosin so that it “lets go” of actin, reducing the excessive connections between them. This action allows your heart muscle to relax more.

Connected myosin

Resting myosin

Actin

CAMZYOS binding to myosin

In people with symptomatic obstructive HCM, reducing the connections between myosin and actin may decrease the obstruction and improve heart function (how the heart relaxes and fills with blood).

To see CAMZYOS in action visit [camzyos.com/about](https://camzyos.com/about)



## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:  
✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.  
It is not known if CAMZYOS is safe and effective in children.

## WARNING: RISK OF HEART FAILURE

! CAMZYOS may cause serious side effects, including heart failure.  
Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS. Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.

! Get medical help right away if you experience new or worsening symptoms, including:  
• shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

The risk of heart failure is also increased when CAMZYOS is taken with certain other medications. Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program. This means:

✓ You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.

✓ Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.

✓ You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.

i If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](https://www.CAMZYOSREMS.com) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.

! CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).

! Get medical help right away if you experience new or worsening symptoms, including:  
• shortness of breath • fatigue • leg swelling  
• chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.

💬 Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.

💬 Talk to your healthcare provider for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.





“My obstructive HCM symptoms left me feeling like a bystander. But now I’ve gone from sitting on the sidelines to being back in the game.”

THIS IS

# MY CAMZYOS MOMENT

*Stephanie, a real CAMZYOS patient*

Individual results may vary.

## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.


## WARNING: RISK OF HEART FAILURE

 CAMZYOS may cause serious side effects, including **heart failure**.

Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS. Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.

-  **Get medical help right away** if you experience new or worsening symptoms, including:
- shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

The risk of heart failure is also increased when CAMZYOS is taken with certain other medications. Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.


Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program. This means:

- ✓ You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.
- ✓ Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.
- ✓ You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.
- i If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?


A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.

 CAMZYOS may cause serious side effects, including **heart failure** (a condition where the heart cannot pump with enough force).


-  **Get medical help right away** if you experience new or worsening symptoms, including:
- shortness of breath • fatigue • leg swelling
  - chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.

-  **Tell your healthcare provider** if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.

-  **Talk to your healthcare provider** for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.



# When compared with placebo

## How was CAMZYOS shown to help in 2 clinical studies?

In the first study, CAMZYOS improved symptoms\* and the ability to be active.†

A **30-week clinical study of 251 people** compared two groups of adults with symptomatic obstructive HCM. One group (123 people) took CAMZYOS while the other group (128 people) took a placebo. All participants who enrolled in the study were symptomatic (73% had NYHA Class II symptoms and 27% had Class III symptoms\*). Though it wasn't a requirement, most of the participants (231) remained on their previous medicine, a beta blocker or calcium channel blocker.

\*Symptoms were measured using the New York Heart Association, or NYHA, classification.  
†The ability to be active was determined by measuring peak oxygen consumption, or pVO<sub>2</sub>, which is the maximum amount of oxygen the body uses during exercise. This was measured during exercise testing on a bicycle or treadmill.  
\*Some improvement=improved pVO<sub>2</sub> by at least 1.5 mL/kg/min.  
More improvement=improved pVO<sub>2</sub> by at least 3.0 mL/kg/min.

### The study first looked at how many people experienced either:



Some improvement‡ in their ability to be active  
**and** an improvement in their symptoms  
  
**or**  
  
More improvement‡ in their ability to be active  
**and** no worsening of their symptoms

Results showed that **more people taking CAMZYOS achieved this outcome** (37%, 45/123) compared with those taking a placebo (17%, 22/128).



**Placebo**—a pill that contains no medicine.  
**NYHA classification**—a method that doctors use to place people into 1 of 4 categories based on their symptoms of heart failure.  
Class I: No symptoms or exercise limitations. Class II: Mild symptoms, some exercise limitations. Class III: Noticeable symptoms with minimal physical activity; comfortable only at rest. Class IV: Severe symptoms while at rest.

CAMZYOS will not work for everyone. Individual results may vary.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Learn more at [camzyos.com](#)

**CAMZYOS**<sup>™</sup>  
(mavacamten)  
2.5, 5, 10, 15mg capsules

### What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:  
✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.  
It is not known if CAMZYOS is safe and effective in children.

### WARNING: RISK OF HEART FAILURE

**CAMZYOS may cause serious side effects, including heart failure.**  
Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

**You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS.** Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.

**Get medical help right away** if you experience new or worsening symptoms, including:  
• shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

**The risk of heart failure is also increased when CAMZYOS is taken with certain other medications.** Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

**Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program.** This means:

- You and your healthcare provider must both be enrolled in the CAMZYOS REMS program** in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.
- Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely**, which includes returning for echos when advised by your healthcare provider.
- You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program.** Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.
- If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](#) or call 1-833-628-7367.

### What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.

**CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).**

**Get medical help right away** if you experience new or worsening symptoms, including:  
• shortness of breath • fatigue • leg swelling  
• chest pain • racing heart (palpitations) • rapid weight gain

### What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.

**Tell your healthcare provider** if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.

**Talk to your healthcare provider** for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.




# When compared with placebo in the first clinical study, CAMZYOS\*:

## Reduced the obstruction


At the beginning and end of the study, the obstruction in people’s hearts was measured with an echocardiogram after they exercised. This was done to find out how much blood was able to pump out of the heart in millimeters of mercury, or mmHg. Doctors call this measurement the left ventricular outflow tract, or LVOT, gradient. A lower LVOT gradient means there is less obstruction in the heart.

 Obstruction to blood flow within the heart decreased **4x more** with CAMZYOS when compared with placebo.

People who took CAMZYOS had an average decrease in their LVOT gradient of 47 mmHg (started at 86 mmHg and went down to 38 mmHg), and people who took a placebo had an average decrease of 10 mmHg (started at 84 mmHg and went down to 73 mmHg).




**LVOT gradient**—When the heart contracts, oxygen-rich blood goes from the heart and out into the body through a pathway called the left ventricular outflow tract, or LVOT. The LVOT gradient is a measurement of blood pressure through this pathway. **A lower LVOT gradient means that more blood is able to flow from the heart.**




## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:  
✔ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.  
It is not known if CAMZYOS is safe and effective in children.

## WARNING: RISK OF HEART FAILURE

 **CAMZYOS may cause serious side effects, including heart failure.**  
**Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.**  
**You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS.** Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.  
People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.

 **Get medical help right away** if you experience new or worsening symptoms, including:  
• shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain


**The risk of heart failure is also increased when CAMZYOS is taken with certain other medications.** Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.


**Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program.** This means:

- ✔ You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.
- ✔ Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.
- ✔ You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.
- i If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?


A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.

 **CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).**


 **Get medical help right away** if you experience new or worsening symptoms, including:  
• shortness of breath • fatigue • leg swelling  
• chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.

 **Tell your healthcare provider** if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.


 **Talk to your healthcare provider** for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Improved symptoms

Symptom improvement was defined as a lowered NYHA class at the end of the study.



~2 out of 3 people taking CAMZYOS had their symptoms improve.

Of those taking CAMZYOS, 65% (80/123) improved by one or more NYHA class compared with 31% (40/128) of those taking placebo.

NYHA classification	Symptoms
Class I	No symptoms or exercise limitations
Class II	Mild symptoms, some exercise limitations
Class III	Noticeable symptoms with minimal physical activity; comfortable only at rest
Class IV	Severe symptoms while at rest

\*This spread covers secondary endpoints.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Learn more at [camzyos.com](#)

Improved quality of life, as reported by patients

How much people were bothered by their symptoms and experienced physical limitations was assessed using the Kansas City Cardiomyopathy Questionnaire (23-item version)—Clinical Summary Score (KCCQ-CSS). Higher scores represent better health status.



3x greater improvement in symptom burden and physical limitations

From when they started the study to when it ended, people taking CAMZYOS reported an average change in score of +14, and those who took a placebo reported an average change in score of +4. The average baseline score was 71 for patients starting the trial.

The frequency and severity of people’s shortness of breath was assessed using the Hypertrophic Cardiomyopathy Symptom Questionnaire—Shortness-of-Breath subscore (HCMSQ-SoB). Lower scores represent less shortness of breath.



3x greater improvement in shortness of breath

From when they started the study to when it ended, people taking CAMZYOS reported an average change in score of -3, and those who took a placebo reported an average change in score of -1. The average baseline score was 5 for patients starting the trial.



What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

WARNING: RISK OF HEART FAILURE



**CAMZYOS may cause serious side effects, including heart failure.**

**Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.**

**You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS.** Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

**The risk of heart failure is also increased when CAMZYOS is taken with certain other medications.** Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

**Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program.** This means:



**You and your healthcare provider must both be enrolled in the CAMZYOS REMS program** in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.



**Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely,** which includes returning for echos when advised by your healthcare provider.



**You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program.** Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.



If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](#) or call 1-833-628-7367.

What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.



**CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).**



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • fatigue • leg swelling • chest pain • racing heart (palpitations) • rapid weight gain

What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.



**Tell your healthcare provider** if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.



**Talk to your healthcare provider** for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.





“I wasn’t always able to keep up with the pace in the city.  
But with fewer symptoms, now I can.”

*Elani, a real CAMZYOS patient, pictured with her mother, Debra*

Individual results may vary.

### What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:  
✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.  
It is not known if CAMZYOS is safe and effective in children.

### WARNING: RISK OF HEART FAILURE

⚠ CAMZYOS may cause serious side effects, including **heart failure**.  
Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS. Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.

📞 ⚠ Get medical help right away if you experience new or worsening symptoms, including:  
• shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

The risk of heart failure is also increased when CAMZYOS is taken with certain other medications. Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program. This means:

- ✓ You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.
- ✓ Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.
- ✓ You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.
- 📄 If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.

### What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.

⚠ CAMZYOS may cause serious side effects, including **heart failure** (a condition where the heart cannot pump with enough force).  
📞 ⚠ Get medical help right away if you experience new or worsening symptoms, including:  
• shortness of breath • fatigue • leg swelling  
• chest pain • racing heart (palpitations) • rapid weight gain

### What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.

💬 Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.

💬 Talk to your healthcare provider for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.



In the second study,

# CAMZYOS reduced the amount of patients who were eligible for, or who chose to undergo, surgery at 16 weeks (avoiding surgery)

A **16-week study of 112 people** compared two groups of adults with symptomatic obstructive HCM. One group (56 people) took CAMZYOS while the other group (56 people) took a placebo. When the study started, everyone had symptoms. Some (7%) had moderate symptoms (NYHA Class II), but most (93%) had severe symptoms (NYHA Class III or IV). Everyone’s obstructive HCM was so advanced that they were eligible for heart surgery.

The study looked at how many people either:

- Remained eligible for surgery at Week 16
- or
- Decided to proceed with surgery prior to or at Week 16

Results showed that **CAMZYOS reduced the proportion of patients who were eligible for, or who chose to undergo, surgery**. 18% of people taking CAMZYOS (~1 out of 5) and 77% of people taking placebo (~4 out of 5) remained eligible for surgery at 16 weeks or chose to undergo surgery prior to or at Week 16. Two people in each group decided to have surgery.



**4 out of 5 people** who took CAMZYOS experienced improvements in their symptomatic obstructive HCM to the point that they **were able to avoid surgery\*** at 16 weeks

vs 1 out of 5 people who took the placebo

\*No longer eligible and did not choose to undergo surgery at Week 16.

Most people in this study (95%) were also taking additional treatments for their oHCM such as a beta blocker, calcium channel blocker, disopyramide, or a combination of these.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Learn more at [camzyos.com](#)



What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

WARNING: RISK OF HEART FAILURE

**CAMZYOS may cause serious side effects, including heart failure.**

Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS. Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.

- Get medical help right away if you experience new or worsening symptoms, including:
  - shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

The risk of heart failure is also increased when CAMZYOS is taken with certain other medications. Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program. This means:

- You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.
- Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.
- You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.
- If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](#) or call 1-833-628-7367.

What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.

**CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).**

- Get medical help right away if you experience new or worsening symptoms, including:
  - shortness of breath • fatigue • leg swelling
  - chest pain • racing heart (palpitations) • rapid weight gain

What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.

- Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.

- Talk to your healthcare provider for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.



# CAMZYOS REMS (Risk Evaluation and Mitigation Strategy)

## Your doctor will help explain your participation in the REMS program.

In some patients, CAMZYOS can cause the heart to become too relaxed, increasing the risk of heart failure. **Because of this risk, CAMZYOS is only available through the CAMZYOS REMS program, which is required by the FDA to monitor the safe use of the medication.**



### Talk to your doctor about how to enroll in the CAMZYOS REMS program.

In order to take CAMZYOS, you must be enrolled in the CAMZYOS REMS program. The program requires you to:

- Review and discuss the **REMS Patient Brochure** with your doctor to understand the risks of CAMZYOS and the symptoms of heart failure, including when to seek medical attention
- Tell your doctor about all prescription and over-the-counter **medicines you take**, including vitamins and herbal supplements, and alert your doctor and pharmacist of any changes to your medications while on treatment. Do not stop or change the dose of a medicine or start a new medicine without telling your doctor
- Get an **echocardiogram** to see if you are eligible for treatment with CAMZYOS (learn more about echocardiograms on page 13)
- Complete the **Patient Enrollment Form** with your doctor
- Get regular **follow-up echocardiograms** at the times recommended by your doctor while on CAMZYOS

If you have questions about the CAMZYOS REMS program, ask your doctor.

Visit [CAMZYOSREMS.com](http://CAMZYOSREMS.com), or call 1-833-628-7367.

In order to prescribe and dispense CAMZYOS, your doctor and pharmacy are required to participate in the CAMZYOS REMS educational program and be certified in the risks and requirements of using CAMZYOS. Because of the special requirements of the CAMZYOS REMS program, CAMZYOS is only dispensed through REMS-certified specialty pharmacies and is not available through local retail pharmacies. Your doctor will give you information on how to get CAMZYOS from a specialty pharmacy. A specialty pharmacy will send CAMZYOS directly to your home and will continue to call you to coordinate medication shipments and help with any medication issues that may arise.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Learn more at [camzyos.com](http://camzyos.com)

## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

## WARNING: RISK OF HEART FAILURE



**CAMZYOS may cause serious side effects, including heart failure.**

Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

**You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS.** Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

**The risk of heart failure is also increased when CAMZYOS is taken with certain other medications.** Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

**Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program.** This means:



**You and your healthcare provider must both be enrolled in the CAMZYOS REMS program** in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.



**Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely**, which includes returning for echos when advised by your healthcare provider.



**You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program.** Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.



If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.



**CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).**



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • fatigue • leg swelling
- chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.



**Tell your healthcare provider** if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.



**Talk to your healthcare provider** for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.



# Monitoring with CAMZYOS


## Regular echocardiograms will help your doctor evaluate how CAMZYOS affects your heart and monitor for safety.

Before you begin CAMZYOS, you will receive an echocardiogram to confirm whether CAMZYOS is right for you. An echocardiogram (also known as an “echo”) is an imaging test, a type of ultrasound, that can be used to look at the heart to see how well it is working.



If you do start on CAMZYOS, you must schedule and attend echocardiograms as instructed by your cardiologist.

These tests allow your doctor to see how your heart is responding to treatment and monitor for safety. Based on echocardiogram results, your doctor may increase, decrease, or keep your dose the same—or may temporarily pause or discontinue your treatment. See echocardiogram schedule below:



Your doctor will let you know if additional echocardiograms are required following any changes or pauses in treatment, or based on echo results.

\*Previously required every 12 weeks.



When you receive your medicine, ask your doctor to schedule your first 3 follow-up echocardiograms. Missed or delayed appointments can lead to an interruption in your treatment.

If you would like to find out about your coverage for these required echocardiograms, please see the back cover for helpful information.

**CAMZYOS**  
(mavacamten)  
2.5, 5, 10, 15mg capsules

## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

## WARNING: RISK OF HEART FAILURE



**CAMZYOS may cause serious side effects, including heart failure.**

**Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.**

You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS. Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

The risk of heart failure is also increased when CAMZYOS is taken with certain other medications. Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program. This means:



You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.



Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.



You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.



If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.



**CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).**



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • fatigue • leg swelling
- chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.



Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.



Talk to your healthcare provider for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.



# Additional precautions around taking CAMZYOS

Your doctor needs to know the answers to these important questions before and during treatment with CAMZYOS:



Are you taking (or do you plan to take, or stop taking) **any medicines other than CAMZYOS?**

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking CAMZYOS with certain medicines or grapefruit juice may cause heart failure.

**Do not stop or change the dose** of a medicine or start a new medicine without telling your healthcare provider.

**Especially tell your healthcare provider if you:**

- Take over-the-counter medicines such as omeprazole (for example, Prilosec®), esomeprazole (for example, Nexium®), or cimetidine (for example, Tagamet®)
- Take other medicines to treat your obstructive HCM disease
- Develop an infection



Do you have **other medical conditions?**  
Are you pregnant or breastfeeding or do you plan to become pregnant or breastfeed?

**CAMZYOS may harm your unborn baby.** Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with CAMZYOS. You may also report your pregnancy by calling the Bristol Myers Squibb reporting line at 1-800-721-5072 or [www.bms.com](http://www.bms.com).

**If you are a female and able to become pregnant:**

- Your healthcare provider will do a pregnancy test before you start treatment with CAMZYOS
- You should use effective birth control (contraception) during treatment with CAMZYOS and for 4 months after your final dose of CAMZYOS
- CAMZYOS may reduce how well some hormonal birth control works. Talk to your healthcare provider about the use of effective forms of birth control during treatment with CAMZYOS

**It is not known if CAMZYOS passes into your breast milk.** Talk to your healthcare provider about the best way to feed your baby during treatment with CAMZYOS.



**Tell your healthcare provider right away if you experience new or worsening shortness of breath, chest pain, fatigue, swelling in your legs, a racing sensation in your heart (palpitations), or rapid weight gain.**

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Learn more at [camzyos.com](http://camzyos.com)



## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

## WARNING: RISK OF HEART FAILURE



**CAMZYOS may cause serious side effects, including heart failure.**

Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

**You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS.** Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

**The risk of heart failure is also increased when CAMZYOS is taken with certain other medications.** Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

**Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program.** This means:



**You and your healthcare provider must both be enrolled in the CAMZYOS REMS program** in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.



**Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely,** which includes returning for echos when advised by your healthcare provider.



**You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program.** Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.



If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.



**CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).**



**Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • fatigue • leg swelling
- chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.



**Tell your healthcare provider** if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.



**Talk to your healthcare provider** for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

# Hear from real patients on CAMZYOS



Stephanie

“Feeling improvement in my symptoms after being on the medicine allowed me to start taking my daily walk again. I’m so glad I can participate in things that I love to do, and now I look forward to doing active things with my husband and with my family.”

—Stephanie



Dani

“Being able to travel—navigate the airport, get on the plane, haul the luggage—that was something I couldn’t have even imagined. So, the fact that I know that I can travel is a welcome change; I know I have the ability to go see my family when I want to. It’s great.”

—Dani



Mike

“I’ve been on CAMZYOS for about a year now, and I’m amazed at what a difference it has made with my symptoms. I can’t do everything I used to, but I’m able to be more active than I’ve been recently.”

—Mike

Individual results may vary.

## More patients on CAMZYOS:



John



Rebecca



Kim



To hear more about what CAMZYOS can do for patients with symptomatic obstructive HCM, visit <https://www.camzyos.com/patient-stories>

These patients were compensated for their time.

## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

## WARNING: RISK OF HEART FAILURE

**!** CAMZYOS may cause serious side effects, including **heart failure**.

Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS. Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.

- !** Get medical help right away if you experience new or worsening symptoms, including:
- shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

The risk of heart failure is also increased when CAMZYOS is taken with certain other medications. Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program. This means:

- ✓ You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.
- ✓ Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.
- ✓ You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.
- i If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.

**!** CAMZYOS may cause serious side effects, including **heart failure** (a condition where the heart cannot pump with enough force).

- !** Get medical help right away if you experience new or worsening symptoms, including:
- shortness of breath • fatigue • leg swelling
  - chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.

- !** Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.

- !** Talk to your healthcare provider for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.



# Interested in sharing your journey?



**YOUR STORY**  
could be just what someone needs to hear...

If you're successfully managing your condition and are being treated with CAMZYOS® what you have to say could **make a positive difference** in another person's life.

Find out how you can “share to inspire” and use your experience for the greater good.



Call us toll-free at **1-855-436-5866** or visit [ShareToInspire.com](https://www.sharetoinspire.com)

Please see accompanying U.S. Full Prescribing Information, including Boxed WARNINGS and Medication Guide for CAMZYOS® in pocket.

 Bristol Myers Squibb®

© 2024 MyoKardia, Inc., a Bristol Myers Squibb company.  
CAMZYOS® and the CAMZYOS and MyCAMZYOS Logos are trademarks of MyoKardia, Inc.  
3500-US-2400030 04/25



Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Learn more at [camzyos.com](https://www.camzyos.com)

**CAMZYOS**  
(mavacamten)  
2.5, 5, 10, 15mg capsules

## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.

## WARNING: RISK OF HEART FAILURE



**CAMZYOS may cause serious side effects, including heart failure.**

Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS. Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.



**Get medical help right away** if you experience new or worsening symptoms, including:

• shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain

The risk of heart failure is also increased when CAMZYOS is taken with certain other medications. Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program. This means:



You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.



Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.



You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.



If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](https://www.camzyosrems.com) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?

A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.



**CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).**



**Get medical help right away** if you experience new or worsening symptoms, including:

• shortness of breath • fatigue • leg swelling  
• chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.



**Tell your healthcare provider** if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.



**Talk to your healthcare provider** for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.





“My obstructive HCM symptoms were slowing me down. But with my symptoms improving, I've gone from taking breaks to taking on tasks on the family farm again.”

THIS IS

# MY CAMZYOS MOMENT

Susan, a real CAMZYOS patient

Individual results may vary.


## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- ✓ adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.


## WARNING: RISK OF HEART FAILURE

 CAMZYOS may cause serious side effects, including **heart failure**.

Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.

You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS. Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.

People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.

 **Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • chest pain • fatigue • racing heart (palpitations) • leg swelling • rapid weight gain


The risk of heart failure is also increased when CAMZYOS is taken with certain other medications. Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.


Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program. This means:

- ✓ You and your healthcare provider must both be enrolled in the CAMZYOS REMS program in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.
- ✓ Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely, which includes returning for echos when advised by your healthcare provider.
- ✓ You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program. Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.
- i If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.

## What are the serious side effects of CAMZYOS?


A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.

 CAMZYOS may cause serious side effects, including **heart failure** (a condition where the heart cannot pump with enough force).


-  **Get medical help right away** if you experience new or worsening symptoms, including:
- shortness of breath • fatigue • leg swelling
  - chest pain • racing heart (palpitations) • rapid weight gain

## What are the most common side effects?

The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.

 Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.

 Talk to your healthcare provider for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.



# Important Facts About CAMZYOS

This is a summary of important information that you need to know about CAMZYOS. Your healthcare provider can work with you to help answer any questions you may have about this medication. Keep this information in a safe place so you can refer to it before and during your treatment.

Look out for the following icons as you read:

Talk to your healthcare provider

Call a healthcare provider right away

Helpful information to remember


## What is CAMZYOS?

CAMZYOS is a prescription medicine used to treat:

- adults with **symptomatic obstructive hypertrophic cardiomyopathy** (obstructive HCM). CAMZYOS may improve your symptoms and your ability to be active.

It is not known if CAMZYOS is safe and effective in children.


## WARNING: RISK OF HEART FAILURE

 CAMZYOS may cause serious side effects, including **heart failure**.

**Heart failure, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death.**

**You must have echocardiograms before you take your first dose and during your treatment with CAMZYOS to help your healthcare provider understand how your heart is responding to CAMZYOS.** Echocardiograms (echos) are imaging tests, a type of ultrasound study, which take pictures of your heart and show how well your heart is working.





People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS.

 **Get medical help right away** if you experience new or worsening symptoms, including:

- shortness of breath • fatigue • leg swelling • chest pain
- racing heart (palpitations) • rapid weight gain

**The risk of heart failure is also increased when CAMZYOS is taken with certain other medications.** Tell your healthcare provider about the medicines you take, both prescribed and obtained over-the-counter, before and during treatment with CAMZYOS.

**Because of the serious risk of heart failure, CAMZYOS is only available through a restricted program called the CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program.** This means:


-  **You and your healthcare provider must both be enrolled in the CAMZYOS REMS program** in order to have it prescribed. Talk to your healthcare provider about how to enroll and for more information about the program.
-  **Your healthcare provider and pharmacist will make sure you understand how to take CAMZYOS safely,** which includes returning for echos when advised by your healthcare provider.
-  **You can get CAMZYOS only from a certified pharmacy that is part of the CAMZYOS REMS program.** Talk to your healthcare provider about finding a certified pharmacy. You will not be able to get CAMZYOS from a local pharmacy.
-  If you have questions about the CAMZYOS REMS Program, ask your healthcare provider, visit [www.CAMZYOSREMS.com](http://www.CAMZYOSREMS.com) or call 1-833-628-7367.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.


Learn more at [camzyos.com](http://camzyos.com)




## What should I discuss with my healthcare provider before starting CAMZYOS?

 Talk to your healthcare provider about **all of your medical conditions**, including if:

 You are **pregnant or plan to become pregnant**—CAMZYOS may harm your unborn baby

 **Tell your healthcare provider right away** if you become pregnant or think you are pregnant during treatment with CAMZYOS. You may also report your pregnancy by calling Bristol Myers Squibb at 1-800-721-5072 or by visiting [www.bms.com](http://www.bms.com).


If you are a female and are able to become pregnant, your healthcare provider will do a pregnancy test before you start taking CAMZYOS. You should use an effective method of birth control (contraception) during treatment and for at least 4 months after your last dose of CAMZYOS.

 CAMZYOS may reduce how well hormonal birth control works. Talk to your healthcare provider about the use of effective forms of birth control during treatment with CAMZYOS.

 You are **breastfeeding or plan to breastfeed**

It is not known if CAMZYOS passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with CAMZYOS.

## What medicines should I discuss with my healthcare provider?


 Tell your healthcare provider about **all of the medicines you take before and during treatment with CAMZYOS**, including:

- prescription medicines
- over-the-counter medicines
- vitamins
- herbal supplements

**Taking CAMZYOS with certain medicines or grapefruit juice may cause heart failure.**

 **Especially tell your healthcare provider if you:**

- take over-the-counter medicines, including medications like omeprazole (for example, Prilosec®), esomeprazole (for example, Nexium®), or cimetidine (for example, Tagamet®)
- take other medicines to treat your obstructive HCM disease
- develop an infection

 **Do not stop or change the dose of a medicine or start a new medicine without telling your healthcare provider.**

Continued on the next page.







# Important Facts About CAMZYOS (continued)

## What are the serious side effects of CAMZYOS?


A **serious side effect** is a side effect that can sometimes become life-threatening and can lead to death.

 **CAMZYOS may cause serious side effects, including heart failure (a condition where the heart cannot pump with enough force).**


-  **Get medical help right away** if you experience new or worsening symptoms, including:
- shortness of breath
  - chest pain
  - fatigue
  - racing heart
  - leg swelling
  - rapid weight gain (palpitations)

## What are the most common side effects?


The most common side effects of CAMZYOS include **dizziness** and **fainting (syncope)**.


 **Tell your healthcare provider** if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of CAMZYOS.

 **Talk to your healthcare provider** for more information about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Bristol Myers Squibb at 1-800-721-5072.

## How should I take CAMZYOS?


 Before starting treatment with CAMZYOS, your healthcare provider will do an echocardiogram (echo) to assess your obstructive HCM. Your heart must be regularly monitored with echos while on treatment to help your healthcare provider understand how your heart is responding to CAMZYOS.

 Talk to your healthcare provider for more information about echos and your echo schedule.

Please see additional Important Safety Information, throughout, including **Boxed WARNING**, [US Full Prescribing Information](#) and [Medication Guide](#) for CAMZYOS.

Learn more at [camzyos.com](#)


## How should I take CAMZYOS? (continued)

 CAMZYOS comes in immediate release capsules, containing 2.5, 5, 10, or 15 mg of mavacamten. Each capsule is marked with the word MAVA and the number of milligrams (mg) it contains.


**Inactive ingredients:** croscarmellose sodium, hypromellose, magnesium stearate (non-bovine), mannitol, and silicon dioxide.  
**Capsule:** contains black iron oxide, gelatin, red iron oxide, titanium dioxide, and yellow iron oxide.

**Take CAMZYOS exactly as your healthcare provider tells you.**


- ✓ 1 time a day, by mouth
- ✗ **Do not** break, open, bite, or chew the capsule
- ✓ Swallow the capsule whole
- ✗ **Do not** change your dose of CAMZYOS without talking to your healthcare provider first

 Talk to your healthcare provider about the right dose for you.

## What if I miss a dose?

- ✓ Take it as soon as possible and take your next dose at your regularly scheduled time the next day.
- ✗ **Do not** take 2 doses on the same day to make up for a missed dose.
-  Your healthcare provider may change your dose, temporarily stop, or permanently stop your treatment with CAMZYOS if you have certain side effects.

## What if I take too much CAMZYOS?

 **Call your healthcare provider or go to the nearest hospital emergency room right away.**

## How should I store CAMZYOS?

Store CAMZYOS capsules between 68°F-77°F (20°C-25°C).  
**Keep CAMZYOS and all medicines out of the reach of children.**



“My obstructive HCM symptoms left me feeling like even the small things were too big. But with CAMZYOS reducing my symptoms, I’ve gone from falling behind to getting back into the swing of things.”

THIS IS  
**MY CAMZYOS  
MOMENT**

Barbara, a real CAMZYOS patient

Individual results may vary.



## Resources for you every step of the way

Once you have your CAMZYOS<sup>®</sup> prescription and enroll in the program, **MyCAMZYOS<sup>®</sup>** is ready to help.

### Patient Access Specialist

Patient Access Specialists will help to:

- Explain your insurance coverage for CAMZYOS
- Understand your out-of-pocket costs
- Inform you about resources that may be available to you

### Free Trial Program

One-time, 35-day free trial for first-time patients

### Co-Pay Program

Commercially insured patients may be eligible to pay as little as \$10 per month for their CAMZYOS prescription.



### Imaging Co-Pay Program

Eligible commercially insured patients may pay as little as \$0 in out-of-pocket costs for required imaging tests, subject to an annual maximum benefit of \$2,500.

Please see Program Terms, Conditions, and Eligibility Criteria in pocket.

**In addition, you have the option to enroll in the MyNurse Navigator Program to receive:**



Important information to help you better understand CAMZYOS and how it works



Tips, tools, and other resources to help you along your treatment journey



Access to a dedicated Nurse Navigator\* who can help answer questions you may have about CAMZYOS<sup>†</sup>

***If you would like to sign up by phone or have any general questions about CAMZYOS, call 855-CAMZYOS (855-226-9967) 8 AM to 8 PM ET, Monday through Friday. You may also sign up online at [MycAMZYOS.com](https://www.MycAMZYOS.com)***

Ask your doctor about enrolling you in **MyCAMZYOS**.

\*Nurse Navigators can provide general information about CAMZYOS but cannot provide medical advice. Your doctor is the best source of information about your health.

<sup>†</sup>Available from 8 AM to 8 PM ET, Monday through Friday. At all other times, nurses will usually return your calls the following business day. Response times may vary in Puerto Rico.