

REIMBURSEMENT GUIDE FOR HEALTHCARE PROVIDERS

MyCAMZYOS® ECHOCARDIOGRAM CO-PAY ASSISTANCE PROGRAM

INDICATION

CAMZYOS® (mavacamten) is indicated for the treatment of adults and pediatric patients weighing 30 kg or more with symptomatic obstructive hypertrophic cardiomyopathy (HCM) to improve functional capacity and symptoms.

IMPORTANT SAFETY INFORMATION

WARNING: RISK OF HEART FAILURE

CAMZYOS reduces left ventricular ejection fraction (LVEF) and can cause heart failure due to systolic dysfunction.

Echocardiogram assessments of LVEF are required prior to and during treatment with CAMZYOS. Initiation of CAMZYOS in patients with LVEF <55% is not recommended. Interrupt CAMZYOS if LVEF is <50% at any visit or if the patient experiences heart failure symptoms or worsening clinical status.

Concomitant use of CAMZYOS with certain cytochrome P450 inhibitors or discontinuation of certain cytochrome P450 inducers may increase the risk of heart failure due to systolic dysfunction; therefore, the use of CAMZYOS is contraindicated with the following:

- Strong CYP2C19 inhibitors
- Moderate to strong CYP2C19 inducers or moderate to strong CYP3A4 inducers

Because of the risk of heart failure due to systolic dysfunction, CAMZYOS is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the CAMZYOS REMS Program.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).



CAMZYOS® Echocardiogram Co-Pay Assistance Program*

This guide is designed to help you understand the **CAMZYOS Echocardiogram Co-Pay Assistance Program** and how to submit for reimbursement on behalf of your patient for their out-of-pocket costs for Risk Evaluation and Mitigation Strategy (REMS) required echocardiograms. The program includes a medical benefit offer for reimbursement of your patient's out-of-pocket costs for required echocardiogram procedures where the full cost is not covered by their insurance.

Patients may pay as little as \$0 in out-of-pocket costs per echocardiogram procedure, subject to an annual maximum benefit of \$2,500. Patients are responsible for any costs that exceed the maximum benefit.

The program does not reimburse for other associated costs such as supplies, office visits, or physician-related services including interpretation of echocardiograms.

Your office or your patient can submit for reimbursement of their echocardiogram procedure out-of-pocket costs after insurance coverage has been applied.

*Please [click here](#) for Program Terms, Conditions, and Eligibility Criteria.



For more information about the program please call
855-CAMZYOS, 8 AM to 8 PM ET, Monday–Friday.

The accurate completion of reimbursement- or coverage-related documentation is the responsibility of the healthcare provider and patient. Bristol Myers Squibb and its agents make no guarantee regarding reimbursement for any service or item.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS™
(mavacamten) capsules
1, 2.5, 5, 10, 15mg

CAMZYOS® Echocardiogram Co-Pay Assistance Program Terms & Conditions

Eligibility Requirements

- Patients must have commercial (private) insurance and must be treated with CAMZYOS for an on-label indication
- Eligible patients must be 18 years of age or older or, if under the age of 18, be enrolled by a parent, legal guardian, or other legally authorized representative
- Patients must live in the United States or United States territories
- Patients are not eligible if they have medical insurance coverage through a state or federal healthcare program, including but not limited to Medicare, Medicaid, Medigap, CHAMPUS, TRICARE, Veterans Affairs (VA), or Department of Defense (DoD) programs; patients who move from commercial plans to state or federal healthcare programs will no longer be eligible
- Patients residing in Massachusetts, Minnesota, or Rhode Island are not eligible

Program Benefits

- The Program includes a medical benefit offer for reimbursement of patient's out-of-pocket costs for required echocardiogram procedures where the full cost is not covered by the patient's insurance; program does not reimburse for other associated costs such as supplies, office visits, or physician-related services, including interpretation of echocardiograms
- Patients may pay as little as \$0 in out-of-pocket costs per echocardiogram procedure, subject to an annual maximum benefit of \$2,500. Patients are responsible for any costs that exceed the maximum benefit
- To receive the Program benefits, a claim must be submitted within 180 days from the date of the Explanation of Benefits (EOB)
- The Program may apply retroactively to out-of-pocket costs for echocardiograms that occurred within 180 days prior to the date of enrollment
- All Program payments are for the benefit of the patient only

Program Timing

- Patients will be evaluated for ongoing eligibility and will continue enrollment in the Program if all eligibility requirements are met

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS™
(mavacamten) capsules 1, 2.5, 5, 10, 15mg

CAMZYOS® Echocardiogram Co-Pay Assistance Program Terms & Conditions (cont'd)

Additional Terms & Conditions

- Patients and prescribers may not seek reimbursement from health insurance, health savings, or flexible spending accounts, or any third party, for any part of the benefit received by the patient through this offer
- Acceptance of this offer confirms that this offer is consistent with patient's insurance. Patients and healthcare providers must report the receipt of co-pay assistance benefits if required by patient's insurance provider
- Offer valid only in the United States and United States territories. Void where prohibited by law, taxed, or restricted
- The Program is not insurance, not transferable, and not conditioned on any past, present, or future purchase
- No membership fees
- Bristol Myers Squibb reserves the right to rescind, revoke, or amend this offer at any time without notice

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS[™]
(mavacamten)^{1, 2.5, 5, 10, 15mg} capsules



Once you have enrolled your patient into the program using the MyCAMZYOS enrollment form, you can submit for reimbursement on your patient's behalf for their echocardiogram procedure out-of-pocket costs after insurance coverage has been applied.

Be sure to have your documentation ready to submit using the online reimbursement portal. You will need to upload copies or a picture of the documentation to your computer or have ready on your mobile device.

Required documentation includes:

- An Explanation of Benefits (EOB) from your patient's insurance company with the following:
 - Date of service
 - CPT code or HCPCS code (echocardiogram procedure code)
 - Patient financial responsibility once insurance has been applied
- CMS-1500 or UB-04 claim form

Claims may be submitted in 3 ways:

1. The online reimbursement portal at www.echocopayportal.com/hcp
2. Fax to: **800-889-2243**
3. Mail to:
Echocardiogram Co-Pay Assistance Program
PO Box 2355
Morristown, NJ 07962

The accurate completion of reimbursement- or coverage-related documentation is the responsibility of the healthcare provider and patient. Bristol Myers Squibb and its agents make no guarantee regarding reimbursement for any service or item.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS™
(mavacamten) capsules
1, 2.5, 5, 10, 15mg

Sample Explanation of Benefits (EOB) Form

EXPLANATION OF BENEFITS - THIS IS NOT A BILL				Please retain this copy for your records and tax purposes					
Member Name: John Sample Member ID #: 12345678-01				Group Name: ABC Company Provider: Smith, Robert			Group #: 123456 Group #: 9876543		
Dates of Service	Service Description	Billed Amount	Allowed Amount	Deductible Amount	Co-pay Amount	Co-insurance Amount	Paid Amount	Amount You Owe	Remark Code
10/14/22	Example Service 99205	\$493.28	\$109.13		\$35		\$74.13	\$35	R
10/5/22	Example Service 99214	\$198.00	\$98.66		\$35		\$63.66	\$35	
Totals		\$691.28	\$207.79		\$70		\$137.79	\$70	

Note: This is a sample form intended for illustrative purposes only.

What are CPT and HCPCS codes and why are they being requested?

CPT and HCPCS codes are used by providers to identify a medical procedure for billing. A CPT code is a 5-digit numerical code that starts with the number 9. An HCPCS code is a 5-digit code that starts with a C followed by 4 numerical digits.



For additional assistance with the online portal or claims submissions, please contact the program at **800-830-1413**, 8 AM to 8 PM ET, Monday–Friday.

NOTE: If the EOB does not include this information, you may submit an itemized receipt of services that includes the information above. Claims should be itemized and should not be bundled.

The accurate completion of reimbursement- or coverage-related documentation is the responsibility of the healthcare provider and patient. Bristol Myers Squibb and its agents make no guarantee regarding reimbursement for any service or item.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS™
(mavacamten) capsules
1, 2.5, 5, 10, 15mg



Instructions for Submitting a Claim in the Portal

Step 1: Visit the online rebate portal at www.echocopayportal.com/hcp

Step 2: Call the program at **800-830-1413** to set up your new account. The program will assist you with creating your account and linking your enrolled patients.

A screenshot of a web portal login page. At the top, it says "Welcome" in bold. Below that, it says "Sign in to your account". There are two input fields: "Email address*" and "Password*". The "Email address*" field has a cursor in it. Below the "Password*" field is a link that says "Forgot password?". At the bottom is a blue button that says "Continue".

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS[™]
(mavacamten) capsules
1, 2.5, 5, 10, 15mg



Step 3: Your dashboard will display a list of your patients. To begin a claims submission, click on **“Submit a Claim”** next to the appropriate patient. If your patient is not listed, please call the program for assistance at **800-830-1413**

Welcome, Marain

Enter 1 or more search criteria to view your eligible patient list below, including patient information and program status. You can also submit new claims for existing patients. If your patient is not listed below, please call us at 800-830-1413.

Search Your Patients

First Name	Last Name	Date of Birth (MM/DD/YYYY) mm/dd/yyyy	Member ID
Product 			
Clear		Search	
Export to Excel			

Patient Name ↑	Date of Birth	Member ID	Product	Enroll Date	Status	Actions
Bms_Uat Patient	03/21/2001	69878618062	Echocardiogram	07/05/2023	Active	Submit a Claim >
Bms_Joseph	03/21/2001	69878581597	Echocardiogram	07/05/2023	Active	Submit a Claim >

The accurate completion of reimbursement- or coverage-related documentation is the responsibility of the healthcare provider and patient. Bristol Myers Squibb and its agents make no guarantee regarding reimbursement for any service or item.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).



Step 4: Click the radio button for the documentation (Claim document/EOB or Other/Misc) that you are uploading. Claim submission will be delayed or rejected if the incorrect button is selected.

Step 5: Upload the required documents by clicking the **“Browse”** button to locate your file.

Step 6: Once all documents are uploaded, click the **“Submit”** button.

NOTE: This page provides important information about the documentation required for your claim to be successfully submitted.

Upload Documents

Please upload the appropriate documents to submit.

Patient

Full Name
Uat Patient Bms

Member ID
69878618062

Document

Please submit the required following documents:

- Claim Form (CMS-1500 or UB-40)
- Patient's Explanation of Benefits (EOB)

OR an itemized receipt from your billing department

Documents need to include all of the following details:

- Facility name
- Patient's name
- Date of service
- Patient financial responsibility once insurance has been applied
- CPT or HCPCS code (echocardiogram procedure code)

All fields are mandatory unless otherwise indicated.

Document Type:

☐ Claim document/EOB

☐ Other/Misc

Information:
You can upload a maximum of 5 files at one time (max size per file is 5MB).
Allowed file types: gif, jpg/jpeg, png, pdf, heic, heif



Browse

or drag it here

Refresh

Submit >

WHAT HAPPENS AFTER A CLAIM IS SUBMITTED?

If you or your patient submits the claim and it is approved, the party that submitted will receive a check in 7-10 business days. The other party will receive an approval notice to keep you both informed. If the claim is not approved, you will both receive a letter informing you of the reason why. You may also receive a call from the program to obtain any missing information to reprocess the claim. If you need assistance at any time, please call **800-830-1413** and ask to speak with a program representative.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS™
(mavacamten) capsules
1, 2.5, 5, 10, 15mg



Additional Features of the Portal

- Create and manage office user accounts for the portal
- Receive notifications of patients' status and communications
- View claims history
- View patient and provider communications history
- View payment reports for your claims
- FAQs about the program
- Resources - Patient Check Request Form for patient's mailed/faxed claims and a sample EOB

Payment Report

From your dashboard within the “**Payment Reports**” tab, you can view payments issued for claims submitted by your office.

Check								
Check Payment Status: Approved: Claim has been approved and awaiting check to be issued Sent to Bank: Funds are pending the bank-to-bank transfer for deposit. Paid: Claim is approved and check has been mailed to the address on file Export to Excel								
Check Transactions								
Product	Patient Name	Member ID	DOS	Payee Info	Amount	Check Confirmation	Status Date	Status
Echocardiogram	Patient at BMS Program	19878399659	08/16/2023	Bms Danbury Townhall Danbury, CA 77777	\$150.00	3343400012	08/16/2023	Paid
Echocardiogram	UAT Patient BMS	69878618062	08/09/2023	Bms Danbury Townhall Danbury, CA 77777	\$100.00		08/16/2023	Sent to Bank
Items per page: 10 1 - 2 of 2								

The accurate completion of reimbursement- or coverage-related documentation is the responsibility of the healthcare provider and patient. Bristol Myers Squibb and its agents make no guarantee regarding reimbursement for any service or item.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).



Provider Profile

Within the **“Profile”** tab, you will be able to make updates to your address, phone number, and fax information.

Provider Profile

Bms Test Profile Details

Practice Name *

Bms Test

Specialty

Address *

Danbury Townhall

Address 2

City *

Danbury

State *

California

Zip Code *

77777

Main Phone Number *

222-222-2222

Fax Number

Email Address *

test@test.com

NPI

8272480137

Tax ID (Optional)

22-2222222

Update >

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS™
(mavacamten) capsules
1, 2.5, 5, 10, 15mg



Claims History

View the status of your patient's claims under the patient profile within the **"Patient Claims History"** tab. You will see if their claim was approved and the amount to be paid, or rejected with the reason for rejection.

Co-Pay Card Details

Print card details

Co-Pay Member ID
69878618062

BIN
000000

PCN
00

Group Number
EX75017001

Submit a Claim >

Patient Profile Details

Patient Claims History

Patient Messages History

Patient Claims History

Product	Claim ID	Claim Type	Date of Service	Amount Paid	Claim Status	Payment Type	Payee
Echocardiogram	1	Medical	08/09/2023	\$100.00	Accepted	Check	Bms Danbury Townhall Danbury, CA 77777

Items per page: 10

1 - 1 of 1

Submit a Claim >



For additional assistance with the online portal or claims submissions, please contact the program at **800-830-1413**, 8 AM to 8 PM ET, Monday–Friday.

The accurate completion of reimbursement- or coverage-related documentation is the responsibility of the healthcare provider and patient. Bristol Myers Squibb and its agents make no guarantee regarding reimbursement for any service or item.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS™
(mavacamten) capsules
1, 2.5, 5, 10, 15mg

IMPORTANT SAFETY INFORMATION (cont'd)

CONTRAINDICATIONS

CAMZYOS is contraindicated with concomitant use of:

- Strong CYP2C19 inhibitors
- Moderate to strong CYP2C19 inducers or moderate to strong CYP3A4 inducers

WARNINGS AND PRECAUTIONS

Heart Failure

CAMZYOS reduces systolic contraction and can cause heart failure or significantly reduce ventricular function. Patients who experience a serious intercurrent illness (eg, serious infection) or arrhythmia (eg, atrial fibrillation or other uncontrolled tachyarrhythmia) are at greater risk of developing systolic dysfunction and heart failure.

Assess the patient's clinical status and LVEF prior to and regularly during treatment and adjust the CAMZYOS dose accordingly. New or worsening arrhythmia, dyspnea, chest pain, fatigue, palpitations, leg edema, or elevations in N-terminal pro-B-type natriuretic peptide (NT-proBNP) may be signs and symptoms of heart failure and should also prompt an evaluation of cardiac function.

Asymptomatic LVEF reduction, intercurrent illnesses, and arrhythmias require additional dosing considerations.

Initiation of CAMZYOS in patients with LVEF <55% is not recommended. Avoid concomitant use of CAMZYOS in patients on disopyramide, ranolazine, verapamil with a beta blocker, or diltiazem with a beta blocker as these medications and combinations increase the risk of left ventricular systolic dysfunction and heart failure symptoms and clinical experience is limited.

CYP450 Drug Interactions Leading to Heart Failure or Loss of Effectiveness

CAMZYOS is primarily metabolized by CYP2C19 and CYP3A4 enzymes. Concomitant use of CAMZYOS and drugs that interact with these enzymes may lead to life-threatening drug interactions such as heart failure or loss of effectiveness.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS™
(mavacamten) capsules
1, 2.5, 5, 10, 15mg



IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS (cont'd)

CYP450 Drug Interactions Leading to Heart Failure or Loss of Effectiveness (cont'd)

Advise patients of the potential for drug interactions, including with over-the-counter medications (such as omeprazole, esomeprazole, or cimetidine). Advise patients to inform their healthcare provider of all concomitant products prior to and during CAMZYOS treatment.

CAMZYOS Risk Evaluation and Mitigation Strategy (REMS) Program

CAMZYOS is only available through a restricted program called the CAMZYOS REMS Program because of the risk of heart failure due to systolic dysfunction. Notable requirements of the CAMZYOS REMS Program include the following:

- Prescribers must be certified by enrolling in the REMS Program
- Patients must enroll in the REMS Program and comply with ongoing monitoring requirements
- Pharmacies must be certified by enrolling in the REMS Program and must only dispense to patients who are authorized to receive CAMZYOS
- Wholesalers and distributors must only distribute to certified pharmacies

Further information is available at www.CAMZYOSREMS.com or by telephone at 1-833-628-7367.

Embryo-Fetal Toxicity

CAMZYOS may cause fetal toxicity when administered to a pregnant female, based on animal studies. Confirm absence of pregnancy in females of reproductive potential prior to treatment and advise patients to use effective contraception during treatment with CAMZYOS and for 4 months after the last dose. Combined hormonal contraceptives (CHCs) containing a combination of ethinyl estradiol and norethindrone may be used with CAMZYOS. However, CAMZYOS may reduce the effectiveness of certain other CHCs. If these CHCs are used, advise patients to add nonhormonal contraception (such as condoms) during concomitant use and for 4 months after the last dose of CAMZYOS.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS[™]
(mavacamten) capsules
1, 2.5, 5, 10, 15mg

IMPORTANT SAFETY INFORMATION (cont'd)

ADVERSE REACTIONS

In the EXPLORER-HCM trial, adverse reactions occurring in >5% of patients and more commonly in the CAMZYOS group than in the placebo group were dizziness (27% vs 18%) and syncope (6% vs 2%). There were no new adverse reactions identified in VALOR-HCM. The safety of CAMZYOS in pediatric patients (≥35 kg) was evaluated in SCOUT-HCM. There were no new adverse reactions compared to those identified in adult studies.

Effects on Systolic Function

In the EXPLORER-HCM trial, mean (SD) resting LVEF was 74% (6) at baseline in both treatment groups. Mean (SD) absolute change from baseline in LVEF was -4% (8) in the CAMZYOS group and 0% (7) in the placebo group over the 30-week treatment period. At Week 38, following an 8-week interruption of trial drug, mean LVEF was similar to baseline for both treatment groups. In the EXPLORER-HCM trial, 7 (6%) patients in the CAMZYOS group and 2 (2%) patients in the placebo group experienced reversible reductions in LVEF <50% (median 48%; range 35-49%) while on treatment. In all 7 patients treated with CAMZYOS, LVEF recovered following interruption of CAMZYOS.

DRUG INTERACTIONS

Potential for Other Drugs to Affect Plasma Concentrations of CAMZYOS

CAMZYOS is primarily metabolized by CYP2C19 and to a lesser extent by CYP3A4 and CYP2C9. Inducers and inhibitors of CYP2C19 and moderate to strong inhibitors or inducers of CYP3A4 may affect the exposures of CAMZYOS.

Impact of Other Drugs on CAMZYOS:

- **Strong CYP2C19 Inhibitors:** Concomitant use increases CAMZYOS exposure, which may increase the risk of heart failure due to systolic dysfunction. Concomitant use is contraindicated
- **Moderate to Strong CYP2C19 Inducers or Moderate to Strong CYP3A4 Inducers:** Concomitant use decreases CAMZYOS exposure, which may reduce CAMZYOS' efficacy. The risk of heart failure due to systolic dysfunction may increase with discontinuation of these inducers as the levels of induced enzyme normalizes. Concomitant use is contraindicated
- **Weak CYP2C19 Inhibitors or Moderate CYP3A4 Inhibitors:** Concomitant use

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS™
(mavacamten) capsules
1, 2.5, 5, 10, 15mg

IMPORTANT SAFETY INFORMATION (cont'd)

DRUG INTERACTIONS (cont'd)

Impact of Other Drugs on CAMZYOS (cont'd)

with a weak CYP2C19 inhibitor or a moderate CYP3A4 inhibitor increases CAMZYOS exposure, which may increase the risk of adverse drug reactions. Initiate CAMZYOS at the recommended starting dosage in patients who are on stable therapy with a weak CYP2C19 inhibitor or a moderate CYP3A4 inhibitor. Reduce dose of CAMZYOS by one level (ie, 15 mg to 10 mg, 10 mg to 5 mg, 5 mg to 2.5 mg, or 2.5 mg to 1 mg [pediatric dosing]) in patients who are on CAMZYOS treatment and intend to initiate a weak CYP2C19 inhibitor or a moderate CYP3A4 inhibitor. Avoid initiation of concomitant weak CYP2C19 and moderate CYP3A4 inhibitors in patients who are on the lowest recommended dose of CAMZYOS. For short-term use (eg, 1 week), interrupt CAMZYOS for the duration of treatment with a weak inhibitor of CYP2C19 or a moderate inhibitor of CYP3A4. CAMZYOS may be reinitiated at the previous dose immediately on discontinuation of concomitant therapy

- Moderate CYP2C19 Inhibitors or Strong CYP3A4 Inhibitors: Concomitant use with a moderate CYP2C19 inhibitor or strong CYP3A4 inhibitor increases CAMZYOS exposure, which may increase the risk of adverse drug reactions. Discontinuing use of a moderate CYP2C19 inhibitor or strong CYP3A4 inhibitor after long-term concomitant use may decrease CAMZYOS exposure, which may reduce CAMZYOS' efficacy. Initiate CAMZYOS at one dose level below the recommended starting dose in patients who are on stable therapy with a moderate CYP2C19 inhibitor or a strong CYP3A4 inhibitor. Reduce dose of CAMZYOS by one level (ie, 15 mg to 10 mg; 10 mg to 5 mg; 5 mg to 2.5 mg; or 2.5 mg to 1 mg [pediatric dosing]) in patients who are on CAMZYOS and intend to initiate a moderate CYP2C19 inhibitor or a strong CYP3A4 inhibitor. Avoid initiation of concomitant moderate CYP2C19 and strong CYP3A4 inhibitors in patients who are on the lowest recommended dose of CAMZYOS. An increase in dose of CAMZYOS may be needed if the moderate inhibitor of CYP2C19 or strong inhibitor of CYP3A4 is discontinued after long-term concomitant use. Monitor for new or worsening symptoms. For short-term use (eg, 1 week), interrupt

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS™
(mavacamten) capsules
1, 2.5, 5, 10, 15mg



IMPORTANT SAFETY INFORMATION (cont'd)

DRUG INTERACTIONS (cont'd)

Impact of Other Drugs on CAMZYOS (cont'd)

CAMZYOS for the duration of treatment with a moderate inhibitor of CYP2C19 or a strong inhibitor of CYP3A4. CAMZYOS may be reinitiated at the previous dose immediately on discontinuation of concomitant therapy

Potential for CAMZYOS to Affect Plasma Concentrations of Other Drugs

CAMZYOS is an inducer of CYP3A4, CYP2C9, and CYP2C19. Concomitant use with CYP3A4, CYP2C9, or CYP2C19 substrates may reduce plasma concentration of these drugs. Closely monitor when CAMZYOS is used with concomitant CYP3A4, CYP2C9, or CYP2C19 substrates unless otherwise recommended in the Prescribing Information.

Certain Combined Hormonal Contraceptives (CHCs): Progestin and ethinyl estradiol are CYP3A4 substrates. Concomitant use of CAMZYOS may decrease exposures of certain progestins, which may lead to contraceptive failure. CHCs containing a combination of ethinyl estradiol and norethindrone may be used with CAMZYOS, but if other CHCs are used, advise patients to add nonhormonal contraception (such as condoms) or use an alternative contraceptive method that is not affected by CYP450 enzyme induction (eg, intrauterine system) during concomitant use and for 4 months after the last dose of CAMZYOS.

Drugs That Reduce Cardiac Contractility

Expect additive negative inotropic effects of CAMZYOS and other drugs that reduce cardiac contractility. Avoid concomitant use of CAMZYOS in patients on disopyramide, ranolazine, verapamil with a beta blocker, or diltiazem with a beta blocker as these medications and combinations increase the risk of left ventricular systolic dysfunction and heart failure symptoms and clinical experience is limited.

If concomitant therapy with a negative inotrope is initiated, or if the dose of a negative inotrope is increased, monitor LVEF closely until stable doses and clinical response have been achieved.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS[™]
(mavacamten) capsules 1, 2.5, 5, 10, 15mg



IMPORTANT SAFETY INFORMATION (cont'd)

SPECIFIC POPULATIONS

Pregnancy

CAMZYOS may cause fetal harm when administered to a pregnant female. Advise pregnant females about the potential risk to the fetus with maternal exposure to CAMZYOS during pregnancy. There is a pregnancy safety study for CAMZYOS. If CAMZYOS is administered during pregnancy, or if a patient becomes pregnant while receiving CAMZYOS or within 4 months after the last dose of CAMZYOS, healthcare providers should report CAMZYOS exposure by contacting Bristol Myers Squibb at 1-800-721-5072 or www.bms.com.

Lactation

The presence of CAMZYOS in human or animal milk, the drug's effects on the breastfed infant, or the effects on milk production are unknown. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for CAMZYOS and any potential adverse effects on the breastfed child from CAMZYOS or from the underlying maternal condition.

Females and Males of Reproductive Potential

Confirm absence of pregnancy in females of reproductive potential prior to initiation of CAMZYOS. Advise females of reproductive potential to use effective contraception during treatment with CAMZYOS and for 4 months after the last dose. CHCs containing a combination of ethinyl estradiol and norethindrone may be used with CAMZYOS. However, CAMZYOS may reduce the effectiveness of certain other CHCs. If these CHCs are used, advise patients to add nonhormonal contraception (such as condoms) or use an alternative contraceptive method during concomitant use and for 4 months after the last dose of CAMZYOS.

Please see additional Important Safety Information on pages 13-18, and full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).

CAMZYOS[™]
(mavacamten) capsules
1, 2.5, 5, 10, 15mg



Supporting Your Patients Every Step of the Way

Have questions or need assistance?



Contact your BMS Access and Reimbursement Manager for general assistance or to schedule a conversation



Call us at **855-CAMZYOS (855-226-9967)**, 8 AM to 8 PM ET, Monday–Friday, to speak with a live Patient Access Specialist



Visit [MyCAMZYOShcp.com](https://www.mycamzyoshcp.com) for information and resources, including the Enrollment Form, to help your patients with access to CAMZYOS

Please see additional Important Safety Information on pages 13-18, and Full Prescribing Information, including **Boxed WARNING**, by [clicking here](#).



© 2026 MyoKardia, Inc., a Bristol Myers Squibb company.
CAMZYOS® and the CAMZYOS and MyCAMZYOS Logos are
trademarks of MyoKardia, Inc.
3500-US-2600589 10/26

CAMZYOS™
(mavacamten) capsules
1, 2.5, 5, 10, 15mg